© Journal of Applied Mathematics & Decision Sciences, 3(2), 189–193 (1999) Reprints Available directly from the Editor. Printed in New Zealand.

Classroom Note

NUMERICAL SOLUTIONS OF NAGUMO'S EQUATION

M. IQBAL

Department of Mathematical Sciences, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia

Abstract. Nagumo's equation is a third order non-linear ordinary differential equation $\frac{d^3u}{dx^3} - c\frac{d^2u}{dx^2} + f'(u)\frac{du}{dx} - (b/c)u = 0$ where f(u) = u(1-u)(u-a), 0 < a < 1. In this paper we have developed a technique to determine those values of the parameters a, b and c which permit

non-constant bounded solutions.

1. INTRODUCTION

Hodgkin and Huxley [11] in their fundamental work on pulses in a squid axon were the first to give a mathematical description of this process. Their model was based on a concept derived from 'Kelvin's Cable Theory' that the nerve membrane is effectively an inductance-free line with a constant capacitance and a non-linear current flow element.

Later a simplified model for the process was proposed by Nagumo, A. Rimoto and Yoshizawa [16] to obtain the non-constant bounded solutions for the third order non-linear ordinary differential equation

$$\frac{d^3u}{dx^3} - c\frac{d^2u}{dx^2} + f'(u)\frac{du}{dx} - (b/c)u = 0$$
(1)

where f(u) = u(1-u)(u-a), 0 < a < 1, and f is a cubic function of u and b is a positive constant, c is the speed of the travelling wave u = u(x + ct). H. Cohen [13], J. Cooley and F. Dodge [4, 5], Green [9], Hagstrom [9] and R. Knight [14] have compiled extensive numerical results for a speed diagram. Many other authors Rinzel [17], Fitzhugh [6] and McKean [15] reviewed the subject for 0 < a < 1, $b \le 0$ and $c \ge 0$.

A natural tool for the mathematical simulation of such processes and phenomena is the theory of impulsive differential equations. At first this theory developed slowly. In the last decade, however, a considerable increase in the number of publications has been observed in various branches of the theory of impulsive differential equations such as Ciment [2].

2. THE ORIGIN, DEVELOPMENT AND SIGNIFICANCE OF NAGU EQUATION

The nervous system consists of nodal points (cell, soma and dendrites), lines (axons) and termini (receptors). The best known aspect of the nervous system is the conduction of the impulse along a single axon. The source of conduction is a pulse which is either rapidly damped out or is shaped into a characteristic form which then propagates down the axon without distortion like a traveling wave.

Hodgkin and Huxley [11, 12, 13], in their fundamental work on pulses in a squid axon were the first to give a successful mathematical description of the process. The Nagumo model is essentially an initial value problem for the following non-linear partial differential equation in the quarter plane $\{u(x,t)|x \ge 0, t \ge 0\}$

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + f - b \int u \, dt \qquad \text{where } f = u(1-u)(u-a), \qquad 0 < a < 1, \quad (2.1)$$

in which u(x,t) is effectively the axon potential and a and b are physiological parameters.

Further to electronically simulate an animal nerve axon, Nagumo and others [16] made an active pulse transmission line using tunnel diodes. This line shapes the signal wave form during transmission, smaller signals are amplified, larger ones are attenuated, narrower ones are widened and those which are wider are shrunk, all approaching the above mentioned wave form.

Differentiating (2.1) with respect to t, we obtain the partial differential equation

$$\frac{\partial 3u}{\partial t \partial x^2} - c \frac{\partial^2 u}{\partial t^2} + f'(u) \frac{\partial u}{\partial t} - (b/c)u = 0.$$
(2.2)

Now if we look for traveling wave solutions u = u(x, t), then on substitution in (2.2) we get the third order O.D.E.

$$\frac{d^3u}{dx^3} - c\frac{d^2u}{dx^2} + f'(u)\frac{du}{dx} - (b/c)u = 0.$$
(2.3)

The parameters a, b and c are to be determined which permit non-constant bounded solutions of (2.3).

3. METHOD OF SOLUTION.

We are dealing with two cases:

(a) When b = 0 then (2.3) reduces to the second order differential equation

$$\frac{d^2u}{dx^2} - c\frac{du}{dx} + f = 0. ag{3.4}$$

(b) When b > 0, then the differential equation (2.3) is reduced to a system of simultaneous equations of the first order by introducing the new variables

$$u' = \frac{du}{dx} = v$$
 and $u'' = \frac{d^2u}{dx^2} = w$.

In vector notations, the system is written as

$${}^{\prime}\vec{u} = \vec{g}(\vec{u}) \quad \text{where } {}^{\prime}\vec{u} = \vec{g}(\vec{u})$$
$${}^{\prime}\vec{u} = \begin{bmatrix} u \\ v \\ w \end{bmatrix} \quad \text{and}$$
$$\vec{g}(\vec{u}) = \begin{bmatrix} v \\ (b/c)u + (3u^2 - 2(1+a)u + a) + cw \end{bmatrix}$$

In solving (3.2) we employed Hamming's [10] predictor corrector method, with Runge-Kutta quartic method to obtain three starting values (points) on the solution curve in addition to the initial point.

4. INITIAL VALUES.

a) For b = 0. The basic requirement now is to obtain initial conditions to get the numerical procedure started. Near u = 0, the linearized equation is

$$\frac{d^2u}{dx^2} - c\frac{du}{dx} - au = 0. \tag{4.1}$$

The auxiliary equation of (4.1) is

$$m^2 - cm - a = 0 (4.2)$$

and has roots

$$m_1 = \frac{c + \sqrt{c^2 + 4a}}{2a}, \ m_2 = \frac{c - \sqrt{c^2 + 4a}}{2a}$$

since parameters a and c are positive, so there is only one positive root $m_1 = \left(\frac{c+\sqrt{b^2+4a}}{2}\right)$. Therefore, the solution $u(x) \simeq Ae^{m_1x}$ and $u'(x) \simeq Am_1e^{m_1x} = m_1u$ for large negative values of $x, u \simeq 0$ and A is constant. Thus the initial conditions used are u(0) = h, $v(0) = u'(0) = m_1h$ where h is a small step size used in the numerical solution and m_1 is the only positive root of (4.2). We used the step size h as h = 0.001 in our calculations.

b) Initial Conditions for b > 0. Near u = 0 the linearized equation is

$$u''' - cu'' - au' - (b/c) = 0.$$
(4.3)

The auxiliary equation is

$$m^{3} - cm^{2} - am - (b/c) = 0. (4.4)$$

Now to obtain the desired behavior of the solution u(x) as $x \to -\infty$, we require that the roots of the cubic (4.4) should be real. Further we also require that only one of the three roots should be positive. These considerations imply that $b < a^2/4$ and c exceeds the largest positive root of the equation

$$(a^2 - 4b)c^4 + 2a(2a^2 - 9b)c^2 - 27b^2 = 0$$
 Burnside [1].

Thus if b and c are chosen satisfying these conditions and m_1 is the only positive root of the auxiliary equation (4.4), then for large negative values of x we must have

$$u \simeq A e^{m_1 x} \text{ for some constant } A$$

$$u' \simeq A m_1 e^{m_1 x} = m_1 u$$

$$u'' \simeq A m_1^2 e^{m_1 x} = m_1^2 u$$

The initial conditions are

$$\begin{array}{c} u = h \\ v = m_1 h \\ w = m_1^2 h \end{array}$$

$$(4.5)$$

where h is the small step size used in the subsequent computations. The value of m_1 was found for the different values of the parameters used by GRAEFFE's ROOT SQUARING procedure (see Froberg [7]).

CONCLUSION.

Our method worked very well and our numerical results supported the conjecture proposed by H.P. McKean [15], that for the value of the parameter a' > 0.5 and b' > 0, no non-constant bounded solution of (4.3) exists.

The underlying idea is that parameter 'a' plays the role of a doping parameter and the disappearance of the non-constant bounded solution corresponds to the physical fact, that if too much of novocaine is injected, the whole nerve goes dead.

Acknowledgments

- 1. The author gratefully acknowledges the excellent research and computer facilities availed at King Fahd University of Petroleum and Minerals during the preparation of this paper.
- 2. The author also gratefully acknowledges the helpful and constructive comments by the editor on the earlier version of the paper.

References

- 1. Burnside and Panton, "Theory of Equations", Vol. 1, Dublin University Press Series (1899).
- 2. Ciment, M. and Leventhal, S.H., "Higher order compact implicit schemes for the wave equation", Math. Computation, Vol. 29(1975), 985-994.
- 3. Cohen, H., "Mathematical developments in Hodgkin Huxley theory and its approximations", Lect. Math. Life Sci., Vol. 8(1976), 89-124.
- Cole, K.S., Antosiewicz, H.A. and Rabinovitz, P., "Automatic computation of nerve excitation", J. Soc. Industrial and Applied Math., Vol. 3(1955), 153-172.
- 5. Dodge, F. and Cooley, J., "IBM Technical Report".
- 6. Fitzhugh, R., "Impulses and physiological states in theoretical models of nerve membrane", Biophysics Journal Vol. 1(1961), 445-466.
- 7. Froberg, C.E., "An Introduction to Numerical Analysis", Addison Wesley Publishing Company (1969).
- Green, M.W. and Sleeman, B.D., "On Fitzhugh's nerve axon equations", J. Math. Biol., Vol. 1(1974), 153-163.
- 9. Hagstrom, T.M. and Keller, H.B., "The numerical calculations of traveling wave solutions of non-linear parabolic equations", SIAM. J. Sci. Stat. Comput. Vol. 7(1986), 978–988.
- Hamming, R.W., "Stable predictor corrector methods for ordinary differential equations", J.A.C.M. Vol. 6(1964), 37-47.
- 11. Hodgkin, A.L. and Huxley, A.F., "A quantitative description of membrane current and its application to conduction and excitation in nerve", Journal of Physiology Vol. 117(1952), 500-544.
- 12. Hodgkin, A.L., "The Conduction of the Nervous Impulse", Liverpool University Press (1971).
- Huxley, A.F., "Ion movements during nerve activity", Ann. N.Y. Acad. Sci. Vol. 81(1959), 221-246.
- Knight, B., "Numerical results for non-linear diffusion systems", Courant Institute of Mathematical Sciences and IBM, T.J. Watson Research Center, Seminar on Partial Differentiation Equations, Summer 1965.
- McKean, Jr. H.P., "Nagumo's equation", J. Advances in Mathematics, Vol. 4 No. 3(1970), 209-223.
- 16. Nagumo, J. Arimoto, S. and Yoshizawa, S., "An active pulse transmission line, simulating nerve axon", Proceedings of the I.R.E., Vol. 50(1962), 2061-2070.
- Rinzel, J., "Hopf bifurcation and repetitive activity under point simulation for a simple Fitzhugh-Nagumo nerve conduction model", J. Math. Biol. Vol. 5(1978), 363.



Advances in **Operations Research**



The Scientific World Journal







Hindawi

Submit your manuscripts at http://www.hindawi.com



Algebra



Journal of Probability and Statistics



International Journal of Differential Equations





Complex Analysis





Mathematical Problems in Engineering



Abstract and Applied Analysis



Discrete Dynamics in Nature and Society



International Journal of Mathematics and Mathematical Sciences





Journal of **Function Spaces**



International Journal of Stochastic Analysis

